

Towards a spatially and temporally constant Karakorum fault slip rate

M.-L. Chevalier^{1*}, J. Van der Woerd², P. Tapponnier³, H. Li¹, F. J. Ryerson⁴, and R. C. Finkel⁵

¹ Institute of Geology, Chinese Academy of Geological Sciences, Beijing, China

² Institut de Physique du Globe de Strasbourg, France

³ Earth Observatory of Singapore

⁴ Lawrence Livermore National Laboratory, USA

⁵ University of California, Berkeley, USA

*mlchevalier@hotmail.com

Detailed study of the Karakorum fault (KF), a major intra-continental strike-slip fault, is important to improve our understanding of the present-day kinematic role of large strike-slip faults in the deformation of Tibet. Indeed, fast rates argue in favor of block models and suggest lateral transfer of material along strike-slip faults [e.g. 1-4], while slow rates argue in favor of continuous deformation models and crustal thickening with minor lateral transfer [e.g. 5]. Recent papers [e.g. 6, 7] suggested that existing velocity fields and geologic rates on active faults in Asia are still too sparse and poorly constrained (both spatially and temporally) to discriminate between a wide range of deformation models. Therefore, new slip-rate data at different timescales and locations are essential to better understand the deformation of Tibet.

The range of geodetic and geologic slip-rates is $\sim 0 - 11$ mm/yr along this structure. At short timescales, InSAR data suggest a rate of 1 ± 3 mm/yr [8] or $0-6 \pm 2$ mm/yr [9], and GPS data yield between $3-4$ mm/yr [10-12] up to ~ 11 mm/yr [13]. At the long-term timescale (Ma), using geochronology, the rate is $\sim 5-10$ mm/yr [e.g. 14-21], even though some authors [22] suggests that the northern section is inactive in the Quaternary.

In order to constrain slip-rates at the Late Quaternary timescale using tectonic-geomorphology, we studied 29 distinct surfaces (8 alluvial surfaces and 8 lateral moraines) located along the southern half of the KF and along its northernmost tip in the Chinese Pamir, that are offset by the fault by varying amounts from 7 to 1520 m. These surface ages were determined using ^{10}Be surface-exposure dating of ~ 300 quartz-rich samples. Our data [23-26] suggest that the KF slip-rate does not decrease toward its tips but is constant along strike, at >5 mm/yr on one fault branch or >7 mm/yr across two branches. This rate is on the same order than that determined by the same technique [4 ± 1 mm/yr on one branch, 27]. In addition to being spatially constant, this late Quaternary rate appears to be, within error, in agreement with most studies at various timescales and suggests that at first approximation, no major discrepancy exists between geodetic and geologic rates.

[1] Armijo, R., Tapponnier, P., Mercier, J.L., and Han, T.-L., 1986. Quaternary extension in southern Tibet: Field observations and tectonic implications. *Journal of Geophysical Research*, 91 : 13,803–13,872.

[2] Peltzer, G., and Tapponnier, P., 1988. Formation and evolution of strike-slip faults, rifts, and basins during the India-Asia collision: An experimental approach. *Journal of Geophysical Research*, 93, 12 : 15,085– 15,117.

[3] Avouac, J.-P., and Tapponnier, P., 1993. Kinematic model of active deformation in central Asia. *Geophysical Research Letters*, 20 : 895–898.

[4] Tapponnier, P., Xu, Z., Roger, F., Meyer, B., Arnaud, N., Wittlinger, G., and Yang, J., 2001. Oblique stepwise rise and growth of the Tibet Plateau. *Science*, 294, 5547 : 1671–1677.

[5] England, P., and Molnar, P., 2005. Late Quaternary to decadal velocity fields in Asia. *Journal of Geophysical Research*, 110, B12401, doi:10.1029/2004JB003541.

[6] Thatcher, W., 2007. Microplate model for the present-day deformation of Tibet. *Journal of Geophysical Research*, Solid Earth, 112, B05411.

[7] Loveless, J. P., and Meade, B. J., 2011. Partitioning of localized and diffuse deformation in the Tibetan Plateau from joint inversions of geologic and geodetic observations. *Earth and Planetary Science Letters*, 303 : 11–24.

[8] Wright, T. J., Parsons, B., England, P. C., and Fielding, E. J., 2004. InSAR Observations of low Slip Rates on the Major faults of Western Tibet. *Science*, 305 (5681): 236-239.

[9] Wang, H., and Wright, T. J., 2012. Satellite geodetic imaging reveals internal deformation of western Tibet. *Geophysical Research Letters*, 39, L07303, doi:10.1029/2012GL051222.

[10] Chen, Q., Freymueller, J.T., Yang, Z., Xu, C., Jiang, W., Wang, Q., and Liu, J., 2004. Spatially variable extension in southern Tibet based on GPS measurements. *Journal of Geophysical Research*, 109, B09401.

[11] Jade, S., Bhatt, B.C., Yang, Z., Bendick, R., Gaur, V.K., Molnar, P., Anand, M.B., and Kumar, D., 2004. GPS measurements from Ladakh Himalaya, India: Preliminary tests of plate-like or continuous deformation in Tibet. *Geological Society of America Bulletin*, 116 : 1385–1391.

[12] Jade, S., Raghavendra Rao, H. J., Vijayan, M. S. M., Gaur, V. K. Bhatt, B. C., Kumar, K., Jaganathan, S., Ananda, M. B., and Dileep Kumar, P., 2010. GPS-derived deformation rates in northwestern Himalaya and Ladakh. *International Journal of Earth Sciences*, doi: 10.1007/s00531-010-0532-3.

- [13] Banerjee, P., and Bürgmann, R., 2002. Convergence across the northwest Himalaya from GPS measurements. *Geophysical Research Letters*, 29 :13.
- [14] Boutonnet, E., Leloup, P. H., Arnaud, N., Paquette, J. L., and Hattori, D. W. J., 2012. Synkinematic magmatism, heterogeneous deformation, and progressive strain localization in a strike-slip shear zone. The case of the right-lateral Karakorum fault. *Tectonics*, doi:10.1029/2011TC003049.
- [15] Lacassin, R., Valli, F., Arnaud, N., Leloup, P.H., Paquette, J.L., Li, H., Tapponnier, P., Chevalier, M.-L., Guillot, S., Maheo, G., and Xu, Z., 2004. Large-scale geometry, offset and kinematic evolution of the Karakorum fault, Tibet. *Earth and Planetary Science Letters*, 219 : 255–269.
- [16] Murphy, M.A., Yin, A., Kapp, P., Harrison, T.M., Ling, D., and Jinghui, G., 2000. Southward propagation of the Karakoram fault system, southwest Tibet: Timing and magnitude of slip. *Geology*, 28 : 451–454.
- [17] Phillips, R.J., Parrish, R.R., and Searle, M.P., 2004. Age constraints on ductile deformation and long-term slip rates along the Karakoram fault zone, Ladakh. *Earth and Planetary Science Letters*, 226 : 305–319.
- [18] Robinson, A. C., 2009a. Geologic offsets across the northern Karakorum fault: Implications for its role and terrane correlations in the western Himalayan-Tibetan orogen. *Earth and Planetary Science Letters*, 279 : 123-130.
- [19] Valli, F., Arnaud, N., Leloup, P. H., Sobel, E. R., Maheo, G., Lacassin, R., Guillot, S., Li, H., Tapponnier, P., and Xu, Z., 2007. Twenty million years of continuous deformation along the Karakorum fault, western Tibet: A thermochronological analysis. *Tectonics*, 26, TC4004, doi:10.1029/2005TC001913.
- [20] Valli, F., Leloup, P.H., Paquette, J.-L., Arnaud, N., Li, H., Tapponnier, P., Lacassin, R., Guillot, S., Liu, D., Deloule, E., Xu, Z., and Maheo, G., 2008. New U-Th/Pb constraints on timing of shearing and long-term slip-rate on the Karakorum fault. *Tectonics*, 27, TC5007, doi:10.1029/2007TC002184.
- [21] Wang, S., Wang, C., Phillips, R. J., Murphy, M. A., Fang, X., and Yue, Y., 2012. Displacement along the Karakoram fault, NW Himalaya, estimated from LA-ICP-MS U–Pb dating of offset geologic markers. *Earth and Planetary Science Letters*, 337-338: 156-163.
- [22] Robinson, A. C., 2009b, Evidence against Quaternary slip on the northern Karakorum Fault suggests kinematic reorganization at the western end of the Himalayan–Tibetan orogen. *Earth and Planetary Science Letters*, 286: 158–170.
- [23] Chevalier, M.-L., Ryerson, F.J., Tapponnier, P., Finkel, R., Van der Woerd, J., Haibing, L., and Qing, L., 2005a. Slip-rate measurements on the Karakorum fault may imply secular variations in fault motion. *Science*, 307, 5708 : 411–414.
- [24] Chevalier, M.-L., Ryerson, F.J., Tapponnier, P., Finkel, R.C., Van Der Woerd, J., Li Haibing, and Liu Qing, 2005b. Response to comment on "Slip-Rate Measurements on the Karakorum fault May Imply Secular Variations in fault Motion". *Science*, 309 : 1326c.
- [25] Chevalier, M.-L., Li, H., Pan, J., Pei, J., Wu, F., Xu, W., Sun, Z., Liu, D., 2011b, Fast slip-rate along the northern end of the Karakorum fault system, western Tibet. *Geophysical Research Letters*, 38, L22309, doi:10.1029/2011GL049921.
- [26] Chevalier, M.-L., Tapponnier, P., Van Der Woerd, J., Ryerson, F.J., Finkel, R.C., and Li, H., 2012. Spatially constant slip rate along the southern segment of the Karakorum fault since 200 ka. *Tectonophysics*, 530-531: 152-179.
- [27] Brown, E.T., Bendick, R., Bourles, D.L., Gaur, V., Molnar, P., Raisbeck, G.M., and Yiou, F., 2002. Slip rates of the Karakorum fault, Ladakh, India, determined using cosmic ray exposure dating of debris flows and moraines. *Journal of Geophysical Research*, 107, B9 : 2192–2205.

Key words: Active Tectonics, Tectonic-Geomorphology, Slip-rate, Karakorum fault, Tibet, Cosmogenic dating