

THE EXPERIMENT IN APPLYING 3D TECHNOLOGY OF MAGNETIC FIELDS INTERPRETATION AT THE ARCHAEOLOGICAL SITE "ARKAIM"

V. KOCHNEV

INSTITUTE OF COMPUTATIONAL MODELING SB RAN,
KRASNOYARSK, RUSSIA
KOCHNEV@ICM.KRASN.RU

G. ZDANOVICH

HISTORICAL CENTER "ARKAIM", CHELYABINSK, RUSSIA
CHELYABINSK@ACES-GROUP.COM

B. PUNEGOV

HISTORICAL CENTER "ARKAIM", CHELYABINSK, RUSSIA
CHELYABINSK@ACES-GROUP.COM

SEE THE CD FOR THE EXTENDED VERSION

ABSTRACT

The technology was elaborated for solving direct and inverse problems of magnetometry. The technology offers the following possibilities:

1. Inverse problems could be solved with the help of anomaly fields got on several layers. The number of levels is not limited.

2. The direct and inverse problems could be solved for a multi layer model. The number of layers is not limited.

The report will reflect the experience in applying this technology for defining the degree of magnet receptivity of the explored space of the settlement Arkaim, the brink of 3d - 2nd millenniums B.C. Maps and sections will be demonstrated and their interpretations will be offered to the audience. Archaeological excavations held on the explored territory gave evidence of the positive results of the offered technologies.

PREREQUISITES OF THE EXPERIMENT

Arkaim, the fortified settlement dated to 1600-1700 B.C., was discovered in 1987 (Arkaim 1995). It is situated 8.2 km north-north-west of Amurskiy, and 2.3 km south-south-east of Alexandronvskiy in the Chelyabinsk region, Russia. "The excavation revealed that the settlement had been burned and, therefore, many details were preserved. The population, however, had vacated the city before the fire and took all their possession with them. Arkaim had two protective circular walls and two circles of standard dwellings separated by a street around a central square. The external wall, 160 m in diameter and 4 m wide, was built from specially selected soil that had been packed into timber frames before being faced with adobe bricks. On the interior, houses abutted the wall

and were situated radially with their doors exiting to the circular internal street" (Koryakova). The site plan is shown at Figure 1.

It is known from experience that the interpretation of magnetic fields and some of their components gives good results in the detection of archaeological objects without actual excavations. The differences in the magnetic susceptibility of the building material of walls, soil and gumus that covered the archaeological object are crucial for the successful detection. That fact was proved by the application of magnetic fields on the archaeological site "Arkaim". We made an attempt to start the construction of the model of the magnetic properties of medium under the surface. For this purpose we used the technology, developed by us for studying different geological objects (Kochnev 1993, 2003).




Figure 1 Archaeological site "Arkaim" (research area is shown in rectangle)




Figure 2 The relief of research area

The software package ADGM-3D is the core element of the technology that makes it possible to calculate magnetic and gravitational fields from known objects generating fields (direct problem), and the parameters of objects from the observed anomalous fields (inverse problem).

THE MODEL OF MEDIUM AND THE METHOD OF THE SOLUTION

For the solution of these problems we assumed a block-layered model with rectangular parallelepipeds as elementary objects. The lateral dimensions of parallelepipeds for the entire

site are the same and are determined by the observation grid. The vertical dimensions are predetermined by the boundaries of layers. We used well-known formulas (Aleksidze 1987:-336) for the solution of the direct problem. For the inverse problem the adaptive method of solving the systems of algebraic (linear and nonlinear) equations (Kochnev 1988:-152, 1993, 1996:120-129) was applied. The method does not accumulate errors and makes it possible to solve the systems with a large number of equations and unknowns quantities.

the second one is situated to the north. (It should be taken into consideration that the anomalies go from South to North almost bias from the bottom left corner to the top right corner) On the profile -1500 we can see a negative anomaly, caused by a natural deepening. Two areas with the calm field (yellow spots in the lower and central parts of the figure) should also be focused at.

SOLUTION OF THE INVERSE PROBLEM

In order to solve the inverse problem it is necessary to assign the initial approximation of model. The solution can be found while applying the problem to the single-layer model, but it proves to be unstable. Minimum discrepancy occurs at the first iteration, but it increases at the iterations following the first one. (Discrepancy is the difference between the original and modeled values of field). After a number of experiments, we accepted the three-layered model of medium with the boundaries of the layers imitating the surface relief. The layers were 30, 60 and 120 cm thick. Thus, the total thickness of model was 210 cm. It was assumed that magnetic susceptibility for all layers is $50 \cdot 10^{-5}$ SI, and the initial approximation error is $10 \cdot 10^{-5}$ SI.




Figure 3 Initial anomalous magnetic field observed at the surface




Figure 4 Modeled magnetic field

INITIAL DATA

Let's move on to the object under study. In Figure 1 we can see the model of "Arkaim" site and the fragment which we used for the experimental interpretation (red lines). The Northwestern segment was excavated earlier, but the eastern part was not, including the area of our research. Figure 2 shows the area relief of the studied fragment. The survey of magnetic field was conducted on the grid of 0.5 x 0.5 m. The dimensions of the studied fragment are 30x40 m. The number of points is $61 \times 81 = 4941$.

The following discrepancies were obtained in the process of solving the inverse problem:

9.7 5.2 3.9 3.1 2.6 nT.

The initial magnetic field (in 10-colors palette) is shown in Figure 3. Negative values are in red colors, positive - in dark-blue. We can see two strips of positive anomalies which coincide with the direction of convex relief forms. The limits of the field vary from -116 to 123 nT. Rectangular forms of slight positive anomalies can be discerned in the background (in dark green) We can also single out two strong anomalies of the complex form. The first is located in the central part (profile 0),




Figure 5 Magnetic susceptibilities for the first layer




Figure 6 Magnetic susceptibilities for the second layer

Archaeological Prospection

In 1 hour we obtained the 3-layer model of magnetic susceptibility for the entire area of study. Figure 4 shows the magnetic field, calculated from the model. It is almost undistinguished from the initial field. The proximity of the initial and the model magnetic fields suggests that one of the possible solutions is obtained.

Note two specific features. The upper layer has high magnetic susceptibility while the lower level is characterized by low magnetic susceptibility. This is especially visible in the area of the decline in the relief.




Figure 7 Magnetic susceptibilities for the third layer. Figure 8 3D model (block variant). Vertical scale is 8 times enlarged. Vertical dimensions in cm, horizontal in meters

ANALYSIS OF THE RESULTS

In Figure 5 we see the magnetic susceptibility of the first layer. It changes from 2 to $91 \cdot 10^{-5}$ SI. At the picture we can distinctly see those special features of the forms which were visible in the magnetic field. Emphasized parts are regions of calm (not technogenic) field.

The distribution of the magnetic susceptibility in the second layer is shown at Figure 6. One can trace here the same features noted before but in a smoother form. The magnetic susceptibility of the third layer (shown here in the individual local palette) we can see in Figure 7. The range is from $13 \cdot 10^{-5}$ to $38 \cdot 10^{-5}$ SI. Figure 8 show the three-dimensional model of magnetic susceptibility. Vertical scale is 8 times enlarged.

Let us further look at the most interesting sections first in the left-to-right, and then in the bottom-to-top direction. On the upper sequence of Figure 9 we see a section along the profile -1500, passing through a negative anomaly, which is located to the south of the bank.

The second sequence presents a section along one of the positive anomalies. Two parts with the increased magnetic susceptibility are noted here: in the center and on the right. Moreover the first one corresponds to the negative form of relief, and the second one - to the positive. The third sequence shows a section along profile +500. In the cavity on the left a reduction in the value of magnetic susceptibility can be seen. Note that the lowered regions of relief both beyond the boundary of

the fortress (sequence 1) and within the limits of the fortress (sequence 3) have lowered values of magnetic susceptibility with smooth changes. Does that indicate that in this part of the fortress there are no walls of dwellings or some other constructions? Could it be a small reservoir inside the fortress, surrounded by trees?

Figure 10 shows profiles that pass from the lower part of the section to the upper. All special features, indicated on the left-to-right profiles, are visible here as well.




Figure 9 SW-NE sections along profiles -1500, 100 and 500

Figure 10 NW-SE sections along lines on 950, 1150, 1550 pickets

THE EXPERIMENTAL PROCESSING OF 4-LEVEL OBSERVATIONS

CONCLUSION

In the small section of 20x1.5 m geophysicists carried out the survey of magnetic field on 4 levels: on the surface, 5, 15 and 25 cm above the surface with a sampling interval 50 cm. The initial and modeled magnetic fields are shown in Figure 11. The upper frame is a section. Let's look at the section and compare them with the results obtained with level 1 of the survey (Fig.9). They somewhat differ.

We obtained a 3-layer, 3-dimensional model of magnetic susceptibility of a near-surface part up to 210 cm in depth (30x40 m) of the archaeological site "Arkaim". This approach enabled us to compose maps of each layer and sections along different paths. We show 6 sections, crossing characteristic anomalous zones. Zones with strong and weak fluctuations of susceptibility were identified on maps and sections.

3 strong anomalies were found. First - negative, beyond the fortress limits, correlated with local lowered form of relief and probably connected with the burial site. Two other ones are positive, inside the fortress, probably connected with ancient metallurgy items.

Data from 4-level survey of 20x1.5 m part were processed. These data turned out to be higher. We would recommend to use this technology advantages for choosing an optimal survey scheme.

The first experiment of the application of this technology showed that there are many interesting tasks in the field of archaeology to be worked at by geophysics and archaeologists in cooperation. We should not forget that magnetometry is a rather precise and highly productive method and at the same time comparatively simple in application.

Topographic survey and mapping are the most laborious part of such works. One of the tasks for archaeologists is to analyze the results of experimental interpretation with the aid of our technology, which was created for solving forward and inverse problems in geology. We worked on this problem with great interest and enthusiasm.




Figure 11 Four curves observations. Plots of initial dt field (up), modeled dt (middle) and section (bottom), visualized block-style




Figure 12 Magnetic fields modeled at profile 100. Plots (up) and section of field modeled on 7 levels

Let's do the calculation of the magnetic field at 7 levels, imitating the surface. The first is on the surface, and the last is at height of 3.5 m. We can see that the magnetic field at latter two levels is close to 0. Analyzing the results of simulation we come to the conclusion that the optimum level of observation could be 0, 15, 30 or 60 cm.

BRIEF DESCRIPTION OF THE TECHNOLOGY

The software package ADGM-3D serves as the basis of the technology. The number of profiles, points, layers and the number of observation levels are unlimited. The rapid and stable method of solving inverse problems in combination with the graphic possibilities gives the possibility to solve complex problems. All illustrations in this paper were obtained with the graphic means of the package.

SPECIAL THANKS

The authors thank the co-authors of the package D.Vasil'ev and V.Sidorov for the collaboration, and also I.Goz., M.Starikova for the help in preparing and translating the work.

REFERENCES

- ALEKSIDZE, M.A., 1987. Approximation methods for solving forward and inverse gravimetric problem. Nauka, Moscow.
- ARKAIM, 1995. Studies and explorations. Chelyabinsk.
- KOCHNEV, V.A., 1988. Adaptive methods of interpreting seismic data. Nauka, Novosibirsk.
- KOCHNEV, V.A., 1993. Adaptive methods of solving inverse geophysical problems. Comp. Center of SB RAN, Krasnoyarsk.
- KOCHNEV, V.A., 1996. Khvostenko V.I - Adaptive methods of solving inverse gravimetric problem. *Geologiya i geofizika* 7:120-129.
- KOCHNEV, V.A., VASILIEV, D.V. and SIDOROV, V.Y., 2003. The technique of solving 3-D gravity problems. SEG International Exhibition, Salt Lake City.
- KORYAKOVA, L. Sintashta-Arkaim Culture (<http://www.csen.org/koryakova2/Korya.Sin.Ark.html>).